


Uva Wellassa University of Sri Lanka
Faculty of Animal Science & Export Agriculture



BASc Degree Programme
Year IV Semester I

End Semester Examination – February 2011/March 2011

ANS 462-2 Scientific writing and project cycle management

Instructions

Answer three (03) questions including question No.1 in booklets provided.

No. of questions : Four (04)
No. of pages : Ten (10)
Time : Two hours (2 hrs)
Total marks allocated : 100%

Index No:

1. Read the research paper given in pages 2 -8 and write an appropriate abstract of **not more than 250 words** .(50 marks)
2. What are the **elements** of a student proposal? Describe what you should include in each of the elements. (25 marks)
3. Describe how **SWOT** analysis helps planning projects? Considering a project on "Development of poultry sector in Uva Province" do a SWOT Analysis and prepare an action plan based on your findings in SWOT. (25 marks)
4. The list of references given in **page 9** is not in alphabetical order. Write the list of references in alphabetical order. If you find any errors in references correct them. (25 marks)

Effect of Force Molting Programs on Egg Production and Quality of Laying Hens

INTRODUCTION

The requirements of the human to animal protein are increasing day by day. Thus more researches are needed to increase production and egg quality. Force molting employed commercially to stop egg production in laying and breeding hens in order to recycle them for another season of egg production. After a molt, the hen's production rate usually peaks slightly below the previous peak rate and egg quality is improved. The point of molting is thus to increase the production, egg quality and profitability of flocks in their second or third laying seasons. Feed withdrawal is the most popular and effective method in molt induction. It has generated several concerns regarding the animal welfare and food safety issues, eventually leading to investigation for alternative approaches (Keshavarz and Quimby, 2002; Webster, 2003; Holt, 2003). Most commercial forced-moulting programmes entail food withdrawal for different periods (Narahari, 2001). McCovan *et al.* (2006) suggest that non-fast induced moulting treatments provide an effective method for inducing moulting in hens and improving their well-being by minimizing discomfort due to food deprivation. The influence of non-feed removal moulting methods on the weight of ovary, oviduct and skeletal quality of hens was examined too (Yildiz and Alpay, 2008). For a complete recovery of the reproductive tract the hen's body weight must drop 30 to 35 percent during the molt. This is achieved by withdrawing the hen's feed completely for as long as two weeks until they go into molt, which means they lose their feathers and cease to lay eggs in addition to losing weight (El-Azim and Ibrahim, 2005). As well as Alodan and Mashaly (1999) reported that the body weight of the moulted hens decreased significantly to 74.5% of their initial body weight and that the hens were returned to a full feed layer ration *ad libitum* and 16 h of light per day (California method) lost 25% of their initial body weight by day 10 of the experiment.

Research is needed to explore the possibility of utilizing locally available protein resources, especially the greens from plants, to replace the conventional protein-rich concentrate meals. It is common practice by farmers in tropical countries to use small amounts of green feed to protect against possible vitamin deficiencies and to provide unidentified growth factors. Further researches are needed to investigate the withdrawing the hen's feed and recovery feeding on different type of diets. Thus the aim of this study was to evaluate the effect of fasting period and recovery feed on different diets as force molting programs on egg production and egg quality traits of laying hens.

MATERIALS AND METHODS

The present study was carried out at the Poultry Farm in the Department of Animal Production, Faculty of Agriculture, South Valley University (Egypt) 2008. Total of one hundred and sixty five 72 week-old white Bovans hens were used in the experiment. All birds divided into four groups (three fasting treatments and control). The first group (control) had thirty hens divided into six replicates of 5 birds each. While, forty-five hens were in each (the second, third and fourth fasting groups). The birds in groups second, third and fourth were fasted for 10, 12 and 15 days, respectively. After fasting periods the groups included three treatments (a crushed yellow corn, crushed yellow corn supplemented with minerals and vitamins and feed layer ration *ad libitum*) each treatment divided into 3 replicates and every replicate included 5 birds (30 birds in control group +45 birds 10 days fasting +45 birds in 12 days fasting +45 birds in 15 days fasting = 160 birds) 5 birds in each replicate were kept in wire cage of 61×55×45 cm in a closed system using

controlled system. The indoor temperature was $20\pm 2^{\circ}\text{C}$ during the experimental period. The control group (natural molting) received 17 h of light/day from 72 weeks of age to the end of the experimental period. The fasting groups received 8 h of light/day during the fasting period and this continued until the end of the three weeks after fasting then increased by half an hour for the first week. After that it was increased one hour per week till the lighting reached 17 h per day and this was continued till the end of the experiment. The composition and calculated analysis of the experimental diet are shown in Table 1. Body weight was recorded before the fasting, the end of the fasting period and end of three weeks after fasting period and after the egg production reached 10% production.

Egg production was recorded daily per hen in each cage during the experimental period and hen day egg production for each group was calculated every week. For each hen, after force molting period feed consumption was recorded and calculated weekly. Feed conversion ratio as g feed/g egg mass was calculated weekly. Egg quality measurements were determined monthly using 9 eggs per group till the end of the experiment. The Haugh unit was measured for the internal quality of the eggs. The height, correlated with the weight, determines the Haugh unit. The higher the number,

Table 1: Composition and calculated analysis of the experimental diets

Ingredients	%
Yellow corn	60.9
Soybean meal (44% Cp)	21.6
Corn gluten meal (60% Cp)	6
Vit and Min. premix*	0.3
Wheat bran	0.45
Dicalcium phosphate	1.36
Calcium carbonate	8.95
Salt	0.4
DL-methionine	0.04
Total	100
Calculated analysis:	
ME (MJ Kg ⁻¹)	11.5
Crude protein (%)	18.4
Crude fiber (%)	2.68
Crude fat (%)	2.78
Ca (%)	3.87
P (Available, %)	0.38
Lysine (%)	0.85
Methionine (%)	0.4

*Vitamins and minerals premix provided per kilogram of the diet: Vit A, 1000 IU; D3 2000 ICU; Vit E, 10 mg; Vit K, 1 mg; B1, 10 mg B2, 5 mg; B6, 1500 mg; B12, 10 mg Pantothenic acid, 10 mg; Nicotinic acid, 30 mg; Folic acid, 1mg; Biotin, 50 mcg; Chloride, 500 mg copper, 10 mg; iron, 50 mg; Manganese, 60 mg; Zinc, 50 mg and Selenium, 0.1 mg

the better the quality of the egg (fresher, higher quality eggs have thicker whites). The test was introduced by Haugh (1937) and is considered the most significant measure of egg quality.

Data were analyzed using a two way Analysis of Variance (ANOVA) to analyze the differences between treatment groups using SAS (1985) General Linear Models procedures. Means were compared using Duncan's multiple range tests. Level of significance used in all results was $p < 0.05$ (Duncan, 1955). The statistical model used for analyzing data obtained was:

$$Y_{ijk} = U + F_i + D_j + (FD)_{ij} + e_{ijk}$$

Where:

- Y_{ijk} = Observed value of the concerned trait
 U = Observed mean for the concerned trait
 F_i = The fixed effect due fasting period
 D_j = The fixed effect due to the type of diet
 $(FD)_{ij}$ = The fixed effect due to the interaction. (Fasting period x type diet)
 e_{ijk} = Random error

RESULTS AND DISCUSSION

Egg production: Effect of fasting periods on egg production values is presented in Table 2. When feed was withheld to induce a force molt, some eggs were laid for several days before egg production dropped to zero. Egg production was reduced to zero by 5th, 4th and 5th days for hens subjected

Table 2: Egg production affected by force molting

Parameter	Fasting days		
	10	12	15
Zero egg production (day)	5	4	5
Beginning of 2nd cycle (first egg) (day)	20	21	25
50% egg production (day)	43	45	57
Peak of egg production (day)	138	150	158

to 10, 12 and 15 days fasting periods, while the number of days out of egg production were 15, 17 and 20 days for hens subjected 10, 12 and 15 days of fasting, respectively. The present results showed that bird reached to 50% egg production at 43, 45 and 57 days after the fasting phase with 10, 12 and 15 days of fasting, respectively. These results means that the 10 days fasting group reached to 50% egg production at 2 and 14 days earlier than the 12 and 15 days group, while the 12 days fasting group reached to 50% egg production at 12 days earlier than the 15 days group. These results are in agreement with the findings of Carey and Brake (1983) who reported that peak of egg production was attained at 138, 150 and 158 days, while the egg production percentage was peaked at 73, 78 and 76%, respectively after deprivation periods in all 10, 12 and 15 days fasting periods in that order. The differences of peak production was 5% between the 10 and 12 days fasting group in favour of 12 days fasting group, while it was found to be 3% between the 10 and 15 days fasting group, in favour of 15 days. The difference obtained for peak of egg production percentago was 2% between the 12 and 15 days fasting group in favour of 12 days fasting group.

Hen Day Egg Production: (HDP) There is insufficient information conducted to evaluate force molting on hen day egg production. The hen day of egg production as affected by the interaction between the recovery feeding and fasting period values are presented in Table 3. The HDP of all fasting groups increased ($p < 0.05$) by about 49, 48 and 46% in groups 2, 3 and 4, respectively as compared with control group. However, no significant differences due to recovery feeding or interaction between fasting period lengths were detected among the four groups. These results are in agreement with the findings of Hassanabadi and Kermanshahi (2007) who found that there were no differences observed between feed removal treatment and nonfeed removal treatment for postmoult hen day production

Egg Weight (EW) and Egg Mass (EM): The Egg Weight (EW) and Egg Mass (EM) as affected by the fasting period and recovery feeding are presented in Table 3. The birds of control group had higher ($p < 0.01$) Egg Weight (EW) by about 2 to 3% than those of all fasting groups but there were no significant differences among fasting treatments. Similar results were obtained by Christmas *et al.* (1985) who indicated that there was no significant difference in egg weight due to fasting treatments of 4 and 10 days. Birds in fasting groups 2, 3 and 4 had higher ($p < 0.01$) egg mass (51, 50 and 48%, respectively) than those of control group. While, non-significant differences were observed in EW and EM due to recovery feeding or interaction between fasting period lengths of the four groups. Hassanabadi and Kermanshahi (2007) who found that there were no differences observed between feed removal treatment and nonfeed removal treatment for postmoult egg weight.

Feed Consumption (FC) and Feed Conversion Ratio (FCR): The FC and FCR as affected by recovery feeding and fasting periods are presented in Table 3. Differences observed in feed consumption due to fasting periods were not significant. The present results indicated that the feed conversion of birds in fasting groups had significant ($p < 0.01$) differences as compared with control group. These results are in agreement with that of Mohamed (1990) who observed no significant differences in feed consumption between 7 and 10 days fasting treatment groups. Hassanabadi and Kermanshahi (2007) who found that there were no differences observed between feed removal treatment and nonfeed removal treatment for feed intake, feed efficiency.

Egg quality: There were no significant differences in egg component percentages due to fasting period length or recovery feeding or fasting period length (Table 4). No significant differences were observed in values of Yolk Percentage (YP) between the eggs of birds fed crushed Yellow Corn (YC), Yellow Corn with Minerals and vitamins (YCM) and Layer Ration (LR) groups. No significant differences were observed for albumen percentage, shell percentage. Also, there were no significant different on the egg quality indices such as egg shape index, shell strength, shell thickness, Haugh units and egg yolk index among treatments. These results due to length of fasting or recovery feeding are presented in Table 5 and 6. The eggshell strength produced by birds fed (LR) was higher ($p < 0.05$) by about 7 and 9% than those eggs produced by hens fed YC and YCM, respectively. Christmas *et al.* (1985) found that forced molting by fasting for 4 and 10 days improved the egg quality. Non-feed removal molting programme based particularly on grain barley

Table 3: Effect of fasting periods and recovery feeding on hen day egg production (HDP)(%), egg weight (gm), Egg Mass (EM), Feed Consumption (FC) and Feed Conversion Ratio (FCR) in white Bovans layer

Items	Hen day egg production (HDP) (%)	Egg weight (gm)	Egg mass (EM)	Feed consumption (FC)	Feed conversion ratio (FCR)
Fasting groups					
Control	32.6±0.74b	61.1±0.4	20.4±0.45b	114.5±0.82a	5.6±0.44a
10 days	48.5±0.87a	59.9±0.80	30.8±0.56a	114.4±0.98b	3.7±0.23b
12 days	47.9±0.08a	59.9±0.67	30.6±0.56a	115.7±0.78a	3.7±0.35b
15 days	47.5±0.80a	50.5±0.81	30.3±0.56a	113.8±0.66a	3.7±0.20b
Probability	0.001	0.817	0.048	0.001	0.005
Recovery feeding					
YC	47.1±0.81	58.8±0.71	30.1±0.53	115.0±0.75	4.1±0.27
YCM	48.0±0.87	59.4±0.80	30.5±0.52	113.3±0.79	3.9±0.26
LR	48.8±0.90	60.3±0.86	31.1±0.58	112.2±0.99	3.9±0.28
Probability	0.403	0.389	0.402	0.142	0.947
Interaction					
10-YC	47.3±1.46	58.0±1.51	29.8±1.05	111.9±2.38	4.3±0.62
10-YC M	48.5±1.52	58.9±1.45	30.8±0.99	108.9±1.20	3.4±0.21
10-LR	49.3±1.35	60.5±1.21	31.8±0.89	113.5±1.20	3.6±0.22
12-YC	47.1±1.55	58.9±1.53	30.2±1.00	118.8±1.24	4.2±0.58
12-YC M	47.8±1.46	60.4±1.22	30.5±0.95	116.8±1.56	4.8±0.74
12-LR	48.7±1.45	60.4±1.20	31.0±0.95	111.6±1.61	4.1±0.55
15-YC	47.0±1.52	59.5±1.45	30.1±0.99	114.2±1.21	3.6±0.20
15-YC M	47.6±1.52	58.9±1.46	30.2±1.01	114.2±1.25	3.5±0.16
15-LR	47.8±1.40	60.1±1.29	30.5±0.92	113.0±1.52	4.1±0.53
Probability	0.987	0.905	0.948	0.5	0.402

a-c Means with different supper scripts in the same column are significantly different (YC) = Crushed yellow corn, (YCM) = Crushed yellow corn with minerals and vitamins, (LR) = Layer Ration

Table 4: Effect of fasting periods and recovery feeding on yolk, shell and albumin percentage in white Bovans layer

Items	Yolk (%)	Shell (%)	Albumin (%)
Fasting groups			
Control	28.0±0.2	12.8±0.1	59.1±0.2
10 days	27.8±0.2	13.6±0.1	59.1±0.3
12 days	28.3±0.3	13.0±0.1	58.6±0.3
15 days	28.7±0.2	13.2±0.1	58.0±0.3
Probability	0.0584	0.6855	0.0776
Recovery feeding			
YC	27.8±0.2	13.2±0.1	58.8±0.3
YCM	28.5±0.2	12.9±0.1	58.4±0.3
LR	28.5±0.2	12.9±0.1	58.5±0.3
Probability	0.1969	0.1871	0.5386
Interaction			
10-YC	27.5±0.4	13.2±0.2	59.2±0.5
10-YC M	27.9±0.5	13.1±0.3	58.8±0.4
10-LR	28.1±0.4	12.5±0.1	59.3±0.5
12-YC	27.5±0.5	12.0±0.2	59.5±0.5
12-YC M	28.9±0.4	13.0±0.3	58.0±0.5
12-LR	27.5±0.5	12.9±0.2	59.5±0.5
15-YC	28.6±0.4	13.9±0.3	57.7±0.7
15-YCM	28.7±0.3	12.7±0.3	58.5±0.5
15-LR	28.8±0.5	13.2±0.2	57.8±0.7
Probability	0.7017	0.032	0.2721

(YC) = Crushed Yellow corn, (YCM) = Crushed yellow corn with minerals and vitamins, (LR) = Layer Ration, Significance level = $p > 0.05$

Table 5: Effect of fasting periods and recovery feeding on shape index, shell strength (Newton) and shell thickness (Micron) in white Bovans layer

Items	Shape index	Shell strength	Shell thickness
Fasting groups			
Control	74.00±0.6	2.6±0.10	380±0.03
10 days fasting	75.00±1.0	2.7±0.9	380±0.03
12 days fasting	73.00±0.9	2.5±0.1	380±0.03
15 days fasting	75.00±0.9	2.5±0.1	380±0.03
Probability	0.146	0.481	0.972
Recovery feeding			
YC	74.00±1.0	2.57±0.1	390±0.03
YCM	75.00±1.0	2.52±0.1	380±0.03
LR	75.00±1.0	2.76±0.1	380±0.03
Probability	0.825	0.331	0.537
Interaction			
10-YC	75.4±1.1	2.6±0.1	390±0.01
10-YCM	75.6±0.7	2.6±0.1	380±0.01
10-LR	74.8±1.0	2.7±0.2	390±0.00
12-YC	71.7±3.2	2.7±0.2	380±0.00
12-YCM	73.8±0.7	2.4±0.2	380±0.01
12-LR	71.7±3.2	2.7±0.2	380±0.00
15-YC	74.7±0.8	2.3±0.1	390±0.01
15-YCM	74.8±0.7	2.4±0.2	390±0.01
15-LR	74.8±1.2	2.8±0.2	380±0.00
Probability	0.797	0.291	0.959

Significance level = $p > 0.05$, (YC) = Crushed yellow corn, (YCM) = Crushed yellow corn with minerals and vitamins, (LR) = layer ration

Table 6: Effect of fasting periods and recovery feeding on yolk index, Haugh unit in white Bovans layer

Items	Yolk index	Haugh unit
Fasting groups		
Control	41.2±0.3	71.2±0.8
10 days fasting	40.9±0.4	71.6±0.9
12 days fasting	40.7±0.4	71.8±0.9
15 days fasting	40.8±0.4	71.0±0.9
Probability	0.906	0.781
Recovery feeding		
YC	40.6±0.4	70.9±0.9
YCM	40.5±0.4	70.6±0.8
L.R.	41.8±0.4	72.4±0.9
Probability	0.35	0.332
Interaction		
10-YC	41.4±0.6	71.1±1.7
10-YCM	40.9±0.6	71.9±1.3
10-LR	40.5±0.8	70.3±1.7
12-YC	40.2±0.7	72.2±1.5
12-YCM	40.0±0.7	71.8±1.5
12-LR	40.2±0.7	72.2±1.5
15-YC	40.3±0.7	69.4±1.8
15-YCM	40.7±0.7	68.3±1.2
15-LR	44.5±0.6	75.3±1.1
Probability	0.343	0.384

(YC) = Crushed yellow corn, (YCM) = Crushed yellow corn, (MYC) = Crushed yellow corn with minerals and vitamins

had positive effect on egg quality traits in laying hens (Petek *et al.*, 2008). From the present result it can be concluded that fasting white Bovans laying hens for 10, 12 or 15 days to induce molting improved performance and egg quality as compared with the natural molting, but recovery feeding after fasting did not have significant effect on hens performance.

CONCLUSION

From the present result it can be concluded that fasting white Bovans laying hens for 10, 12 or 15 days to induce molting improved performance and egg quality as compared with the natural molting, but recovery feeding after fasting did not have significant effect on hens performance.

In this study, fasting groups were the best between control groups. These fasting groups of hens had significantly higher egg production and egg mass when compared with other group. More studies still needed to improve egg production and egg quality in laying hens.

References

- Kubikova, I., Vyboh, P., and Kostal, L. (2001) Behavioural, endocrine and metabolic effects of food restriction in broiler breeder hens. *Acta vet. Brno*. 70:247-257.
- Hocking, P.M., Gilbert, A. B., Walker, M., and Waddington, D. (1987) Ovarian follicular structure of White Leghorns fed ad libitum and dwarf and normal broiler breeders fed ad libitum or restricted until point of lay. *Br. Poult. Sci.* 28:493-506.
- Dewil, E., Darras V.M., Spencer G.S.G., Lauterio T.J., Decuypere, and E. (1999) The regulation of GH dependent hormones and enzymes after feed restriction in dwarf and control chickens. *Life Sci.* 64: 1359-1371
- Hocking, P.M., Waddington, D., Walker, M.A., and Gilbert, A. (1989) Control of the development of the ovarian follicular hierarchy in broiler breeder pullets by food restriction during rearing. *Br. Poult. Sci.* 30:161-174.
- FAWC, (1998) Report on the welfare of broiler breeders. Farm Animals Welfare Council (FAWC), Sorption, Surrey, U.K.
- Fattori, T. R., H. R. Wilson, R. H. Harms, and Miles, R. D. (1991) Response of broiler breeder females to feed restriction below recommended levels. 1. Growth and reproductive performance. *Poultry Sci.* 70:26-36.
- Bennett, C.D., and Lesson, S. (1989) Growth of broiler breeder pullets with skip-a-day versus daily feeding. *Poult. Sci.* 68: 836-838.
- Davison, T.F., Rea, J., and Rowell, J.G. (1983) Effects of dietary corticosterone on the growth and metabolism of immature *Gallus domesticus*. *Gen. Comp. Endocrinol.* 50:463-468.
- Hocking, P. M., (1993) Effects of body weight at sexual maturity and the degree and age of restriction during rearing on the ovarian follicular hierarchy of broiler breeder females. *Br. Poult. Sci.* 34:793-801
- Bartov, I., Bornstein, S., Lev, Y., Pines, M., and Rosenberg, J. (1988) Feed restriction in broiler breeder pullets, skip-a-day versus skip-two-day. *Poult. Sci.* 67: 809-813.
- Hocking, P.M., Maxwell, M.H. and Mitchell, M.A. (1996) Relationships between the degree of food restriction and welfare indices in broiler breeder females. *Br. Poult. Sci.* 37 :263-278.
- Hocking, P.M., Gilbert, A. B., Walker, M., and Waddington, D. (1987) Ovarian follicular structure of White Leghorns fed ad libitum and dwarf and normal broiler breeders fed ad libitum or restricted until point of lay. *Br. Poult. Sci.* 28:493-506.
- Hocking, P.M., Waddington, D., Walker, M.A., and Gilbert, A. (1989) Control of the development of the ovarian follicular hierarchy in broiler breeder pullets by food restriction during rearing. *Br. Poult. Sci.* 30:161-174.
- Katanbaf, M. N., E. A. Dunnington, and Siegel, P. B. (1989) Restricted feeding in early and late-feathering chickens. 2. Reproductive responses. *Poultry Sci.* 68:352-358.