

**ELECTROPHYSIOLOGICAL AND BEHAVIOURAL
RESPONSES OF RED PALM WEEVIL TO SELECTED
PALM VOLATILES**

A dissertation submitted to the
Faculty of Animal Science and Export Agriculture
UvaWellassa University

In partial fulfillment of the requirements for the award of
Bachelor of Science in Palm & Latex Technology and Value Addition

By
UDADENIYE VIYANNALAGE ACHINI PRAMUDIKA

**Palm & Latex Technology and Value Addition Degree
Programme**

**Faculty of Animal Science and Export Agriculture
UvaWellassa University of Sri Lanka**

2018

ABSTRACT

Palm weevil (RPW) *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) is a devastating pest of coconut and other palms species in Sri Lanka and other countries. Being the conceal habitat of the pest inside the palm trunk; management of the pest is difficult and prevention of the pest damage is more important. Pheromone trap is one strategy popularizing among farmers as green pest management method. Increasing the efficiency of aggregation pheromone is a challenge. Use of responsive host volatiles for pest management is a recent trend in the field of pest management. Both electrophysiological and behavioural studies were carried out to select and formulate synergistic semiochemical formulation from six host volatiles and RPW aggregation pheromone. Electroantennogramme (EAG) test results indicated, the highest EAG response (Female: -0.8 ± 0.05 , Male: -0.46 ± 0.09 mV) to pentanol from both female and male. It is always higher than RPW aggregation pheromone (Female: -0.55 ± 0.04 , Male: -0.45 ± 0.04 mV) and other tested five host volatiles. Behavioural studies indicated that, pentanol is an attractive volatile and $81 \pm 3.74\%$ attracts towards the pentanol applied coconut fronds than hexane applied fronds. Further, dose response of EAG, EAG and behaviour studies indicated that, pentanol: pheromone at 1:1 mixture gave the highest response and attraction for both female and male (Male: -0.86 ± 0.27 , Female: -0.511 ± 0.165 mV)- among the tested mixtures of 1:1, 1:2, 1:3, 1:4, 1:5, pheromone alone and pentanol alone. Therefore, pentanol: pheromone at 1:1 formulation can be recommended for use as a semiochemical lure for RPW mass trapping after conducting mass trapping experiments in the field.

Keywords: Electroantennogramme, Host plant volatiles, Mass trapping, Pheromone synergist, Red palm weevil