

**GROWTH PERFORMANCE OF *Holothuria scabra* (SAND FISH)
WITH DIFFERENT STOCKING DENSITY IN OPEN SEA PEN
CULTURE IN JAFFNA, SRI LANKA**

MAHISHALINI VARATHARAJAH

UWU/EX/14/0069

Aquatic Resources Technology Degree Programme

Department of Animal Science

Faculty of Animal Science and Export Agriculture

Uva Wellassa University of Sri Lanka

2018

Abstract

Sea cucumber, *Holothuria scabra* is over exploited species in worldwide due to the high fishing intensity from the wild stocks. Sea cucumber pen culture practices are being expanded in Northern Province of Sri Lanka and presently operated more than 57 pen culture farms in the region. This research is intended to investigate the growth performance of *H. scabra* pen culture with different stocking densities to find out the most appropriate stocking density to introduce the farmers. This research was carried out in coast of Mankumpan village in Jaffna district, Sri Lanka. Early matured *H. Scabra* with average weight 49.78 g and average length 11.28 cm were collected from the Sea cucumber farmers. Four pens were constructed by using 5 mm mesh size high density polyethylene nets and wooden planks. The size of each pen was 25 m². Early matured *H. scabra* was reared with stocking densities of 1, 2, 3 and 4 individual m². Thirty percentage sample size was randomly collected in each pens to measure the growth parameter. Average length, weight and water quality parameters were measured twice a month and finally, survival rate was calculated. Specific Growth Rate at twice a month interval was calculated and it was recorded as $2.53 \pm 0.56 \text{ gday}^{-1}$, $1.98 \pm 0.20 \text{ gday}^{-1}$, $1.95 \pm 0.21 \text{ gday}^{-1}$ and $1.08 \pm 0.27 \text{ gday}^{-1}$ for pen 1, pen 2, pen 3 and pen 4 respectively. Data was analyzed using Minitab17 version by one way ANOVA test. Average survival rates for pen 1 and pen 2 were recorded as 100% and for pen 3 and pen 4 were recorded as 85.33% and 85% respectively. Temperature, Salinity, P^H, Nitrate and Phosphate were ranged 27–31°C, 35–40 ppt, 7.3–8.8, 0.25–2.5 mgL⁻¹ and 0.38–0.44 mgL⁻¹ respectively. The results show that, there is a significant difference ($p < 0.05$) between Specific growth rate and stocking density. The result indicated that the specific growth rate and survival rate were higher in pen 1 and pen 2 than pen 3 and pen 4 and applicable for the implementation.

Keywords: *Holothuria scabra*, Pen Culture, Growth performance, Stocking density